

For adults with plaque psoriasis

Are you ready for relief with DUOBRII Lotion?

Using DUOBRII Lotion is simple. Just remember these steps:



Apply just once a day If possible, apply at night



Apply only to your plaquesBe careful to avoid surrounding, healthy skin



Try not to use too much

You only need a thin layer on each plaque



Use only until your psoriasis is under control Stop using once your plaques have cleared, and tell your doctor if you experience side effects



Helpful tips to keep in mind

- Try to avoid intense sunlight and always wear sunscreen. Your skin will be more sensitive to the sun during treatment
- Don't use moisturizers within an hour of applying

- Check to make sure other skin products you use are not drying out your skin
- Don't apply to skin that is sunburned, infected, or has eczema
- Make sure skin is completely dry when applying

Always talk to your doctor about your experiences and concerns throughout treatment.

If you have more questions during treatment, visit DUOBRII.com

What is DUOBRII® Lotion?

DUOBRII (halobetasol propionate and tazarotene) Lotion, 0.01%/0.045%, is a prescription medicine used on the skin (topical) to treat adults with plaque psoriasis. It is not known if DUOBRII Lotion is safe and effective in children.

Important Safety Information

• DUOBRII Lotion is for use on the skin only; do not use it in your mouth, eyes, or vagina.

DUOBRII Lotion may cause birth defects if used during pregnancy.

Please see additional Important Safety Information on page 2 and click <u>here</u> for full Prescribing Information, including Patient Information.

Duobrii® (halobetasol propionate and tazarotene) Lotion 0.01%/0.045%

Important Safety Information (cont'd)

What is the most important Information I should know about DUOBRII Lotion?

DUOBRII Lotion may cause birth defects if used during pregnancy.

A negative pregnancy test must be obtained before females of child-bearing age start using DUOBRII Lotion and they must use effective birth control during treatment. Begin treatment during a normal menstrual period.

Stop using DUOBRII Lotion and tell your healthcare provider right away if you become pregnant while using DUOBRII Lotion.

Before you use DUOBRII Lotion, tell your healthcare provider if you:

- have eczema or any other skin problems, including skin infections, which may need to be treated before using DUOBRII.
- have diabetes, adrenal gland problems or liver problems.
- are breastfeeding or plan to breastfeed. If you use DUOBRII and breastfeed, do not apply DUOBRII to your nipple area.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

- Especially tell your healthcare provider if you take corticosteroids by mouth or injection or use other skin products that contain corticosteroids.
- Ask your healthcare provider for a list of medicines that may make your skin more sensitive to sunlight.

What should I avoid during treatment with DUOBRII?

- To avoid a severe sunburn, avoid sunlight, including sunlamps and tanning beds, as much as possible, and use sunscreen, protective clothing and a hat while in sunlight. Talk to your healthcare provider if you get sunburn, and do not use DUOBRII Lotion until your sunburn is healed.
- Avoid using DUOBRII on skin with eczema because it may cause severe irritation.

DUOBRII may cause side effects, including:

- If too much DUOBRII passes through your skin it can cause adrenal glands to stop working
- Cushing's syndrome, a condition from too much exposure to the hormone cortisol



- High blood sugar (hyperglycemia)
- · Effects of growth and weight in children
- Skin irritation. If you get too much skin irritation at the site of application, your healthcare provider may tell you to interrupt or stop using DUOBRII or to use it less often.
- Vision problems, including an increased chance of developing cataracts and glaucoma. Tell your healthcare provider about any vision problems during treatment.

The most common side effects of DUOBRII Lotion include: redness, itching, swelling, burning, stinging, application site pain, inflamed hair follicles (folliculitis), thinning of the skin (atrophy), peeling and rash.

To report SUSPECTED ADVERSE REACTIONS, contact Ortho Dermatologics at 1-800-321-4576 or FDA at 1-800-FDA-1088 or visit www.fda.gov/medwatch.

Please click <u>here</u> for full Prescribing Information, including Patient Information.

Ortho Dermatologics

®/TM are trademarks of Ortho Dermatologics' affiliated entities.
© 2020 Ortho Dermatologics' affiliated entities. DUO.0118.USA.20